

REVIEW

Functional Assessment Scales in a General Intensive Care Unit. A Review

Anna Christakou, PT, MSc, PhD, Emmanuel Papadopoulos, PT, MSc, PhD, Irimi Patsaki, PT, MSc, Giorgos Sidiras, RN, MSc, Srafeim Nanas, MD

Critical Care Medicine Department, Evagelismos General Hospital of Athens, University of Athens, Athens, Greece

KEY WORDS: *functional ability; disability; impairment; scales; assessment; intensive care patients*

ABBREVIATIONS

ICU = intensive care unit
 BI = Barthel index
 FIM = functional independence measure
 FSS-ICU = functional status score for the ICU
 PFIT = physical function ICU test
 MRS = modified Rankin scale
 GOS = Glasgow outcome scale
 DRS = disability rating scale

Correspondence to:

Anna Christakou, MSc, PhD,
 Critical Care Medicine Department,
 Evagelismos Hospital,
 Athens, Greece;
 Tel.: +30-6977370284;
 E-mail: achristakou@phed.uoa.gr

*Manuscript received March 5, 2013;
 Revised manuscript received May 24,
 2013; Accepted June 9, 2013*

Conflict of Interest: none declared

ABSTRACT

Patients in the intensive care unit (ICU) are often exposed to prolonged immobilization, thus they lose their functional ability. Therefore, it is crucial to assess patients' functional ability during their stay in and upon their discharge from the ICU. Several scales have been used so far for the assessment of functional ability, impairment and/or disability in ICU patients. These outcome measures include several assessment scales, such as the Barthel Index, the Functional Independence Measure, the Functional Status Score for the ICU, the Physical Function ICU Test Modified Rankin Scale, the Karnofsky Scale Index, the 4P questionnaire, the Glasgow Outcome Scale, and the Disability Rating Scale. The choice of the most appropriate assessment scale will depend on the specific patient population, the diagnosis and rehabilitation phase and the psychological properties of the available measurement. The aim of the present review is to describe the functional assessment scales for ICU, to examine the psychometric evidence for reliability and validity and to summarize the strengths and the weaknesses of these scales.

INTRODUCTION

During an intensive care unit (ICU) stay, patients are often exposed to prolonged bed rest, dysfunction of vital organs, sepsis, hypoxemia and neuromuscular drug toxicity. As a result, the cardiovascular system status may be impaired and critical illness neuromuscular syndromes may occur.¹ Both of these conditions may delay ventilator weaning and increase ICU and hospital stay.² In particular, the prolonged immobility and inactivity may result in loss of muscle strength and endurance, and loss of balance and neuromuscular coordination, further leading to total functional impairment, and thus impaired quality of life. Research has shown that after 1 week of bed rest, muscle strength may decrease as much as 20%, with an additional 20% loss of remaining strength occurring each subsequent week.³ Therefore, examining the functional ability and starting early mobilization of ICU patients should increase the weaning success rate, decrease ICU and hospital stay, and improve their quality of life in the ICU and beyond.^{4,5}

Due to the functional impairment of ICU patients during their stay in the unit, there is a need to assess functional ability upon their discharge from the unit. Furthermore, the cost of caring for the survivors of ICU after their discharge from the unit is quite high and the impact of an effective functional treatment for them would be significant economically and socially. Thus, the development and use of a functional

outcome measure is necessary in order to assess and improve their functional ability, increasing the number of patients who are discharged from ICU. However, the choice of a functional assessment outcome measure is crucial for the evaluation and choice of the best provision of their rehabilitation. Thus, selecting an inappropriate functional assessment outcome measure may lead to incorrect conclusions regarding the benefits of treatment.⁶

Few outcome measures have been used to assess the functional ability of ICU patients.⁷⁻¹⁰ Furthermore, Nickol et al¹¹ and Shukla et al¹² in their reviews discussed general outcome measures for only traumatic brain injury (not in an ICU setting), including quality of life measures. Also, Black et al¹³ describe generic and disease-specific outcome measures of adult critical care survivors. More recently, Gosselink et al¹⁴ reviewed outcome measures for assessing sedation, level of cooperation, cardiorespiratory status, muscle and respiratory strength, and functional performance of ICU patients. There has not been any recent review examining the use and psychometric properties for only functional assessment scales in a general ICU setting, which includes patients with different pathologies, e.g., traumatic brain injuries, general surgeries, respiratory diseases, neuromuscular diseases, etc. Therefore, the aims of the present study were (a) to describe the functional outcome measures of ICU patients, i.e., purpose, number of items, description of subscales, response format and scoring, (b) to examine the psychometric evidence for reliability (internal consistency and test-retest) and validity (content, construct, criterion and discriminant), and (c) to summarize the strengths and the weaknesses of the existing functional outcome measures.

**CHARACTERISTICS
OF A MEASUREMENT SCALE
OF INTENSIVE CARE PATIENTS**

It is useful to determine the characteristics of an assessment scale of intensive care patients. In order for an assessment tool to be used, it should be reliable, reproducible, validated, easy to use and sensitive to any clinical change. Firstly, the scale should be simple to administer to the patient population. The patients' responses should be scored only by the scale instructions and not subjectively. Second, the scale must be reliable which means that it should contain three measurement properties: internal consistency, test-retest reliability and measurement error. Reliability refers to the degree to which the measurement instrument is free from measurement error¹⁵ and estimates the extent to which scores for patients who have not changed are the same for repeated measurements under several conditions: (a) using different sets of items from the same measurement tool (internal-consistency), (b) across time (test-retest), (c) by different persons (i.e., raters) on the same occasion (inter-rater) and (d) by the same person (i.e., raters or responders) on differ-

ent occasions (intra-rater).¹⁶ Third, the scale must be validated, which means that the instrument measures the construct(s) it purports to measure.¹⁵ Validity contains three measurement properties: content validity, construct validity and criterion validity. Fourth, the results of the scale should be reproducible over time. It should have sensitivity and responsiveness to detect small changes which impact function and not demonstrate a ceiling effect as functional ability improves, allowing discrimination between high functioning survivors. Lastly, the scale should ideally be free to administer (i.e., no copyright fee).¹¹

**DESCRIPTION OF THE FUNCTIONAL
ASSESSMENT SCALES IN AN ICU**

The most common scales in the literature which have been used in a general ICU are the following: the Barthel Index, the Functional Independence Measure, the Functional Status Score for the ICU, the Physical Function ICU Test Modified Rankin Scale, the Karnofsky Scale Index, the 4P questionnaire, the Glasgow Outcome Scale, and the Disability Rating Scale (Table 1).

(A) BARTHEL INDEX (BI)

The Barthel Index (BI)¹⁷ has been used to measure physical functioning and was improved upon by the Functional Independence Measure (FIM). It measures the capacity to perform 10 basic activities of daily living. In particular, items are divided into groups that relate to self-care (feeding, grooming, bathing, dressing, bowel and bladder care and toilet use) and mobility (ambulation, transfers and stairs climbing). It gives a quantitative estimation of the patient's level of dependency with scoring from 0 (totally dependent) to 100 (totally independent). The range of BI was described by classifying the patients as having minimal or no disability (BI score, >90), moderate disability (BI score, 55–90) or severe disability (BI score <55). This tool has been reported to have high reliability.¹⁸

(B) FUNCTIONAL INDEPENDENCE MEASURE (FIM)

The FIM¹⁹ is the most widely accepted functional assessment tool used for assessing basic functional activities of ICU patients and their progress during in-patient rehabilitation.^{20,21} Two separate domains of items comprise the motor domain consisting of 13 items and the cognitive domain consisting of 5 items. The FIM is a multi-dimensional measure which assesses self-care, sphincter control, transfers, locomotion, communication, and social cognition in addition to cognitive and motor sub-scales. FIM scores range from 1 to 7; a FIM item score of 7 is categorized as "complete independence," while a score of 1 is "total assistance". Each dimension is then added, yielding a possible total score between 18 (complete dependence) and 126 (complete independence). Precision, inter-rater reliability and validity have been established.²²⁻²⁴ At

TABLE 1. Functional Assessment Scales in the ICU

<p>1) Barthel Index (BI): it measures the capacity to perform 10 basic activities of daily living</p> <ul style="list-style-type: none"> ● self-care (feeding, grooming, bathing, dressing, bowel & bladder care & toilet use) & mobility (ambulation, transfers and stairs climbing) ● scoring ranges from 0 (totally dependent) to 100 (totally independent) ● BI index score >90: minimal or no disability ● BI index 55-90: moderate disability ● BI index <55: severe disability
<p>2) Functional Independence Measure (FIM): the most widely employed functional assessment tool</p> <ul style="list-style-type: none"> ● motor domain (13 items) ● cognitive domain (5 items) ● FIM scores: 1 (total assistance) - 7 (complete independence) for each variable ● FIM total score: 18 (complete dependence) - 126 (complete independence)
<p>3) Functional Status Score for the ICU (FSS-ICU): consists of 3 pre-ambulation categories (rolling; supine to sit transfer; & unsupported sitting) and 2 ambulation categories (sit to stand transfers; & ambulation)</p> <ul style="list-style-type: none"> ● Rating: 1 (total dependent assistance) - 7 (complete independence) scale ● Score: 0-35 (0 score: unable to perform a task due to physical limitations or medical status)
<p>4) 4P questionnaire: evaluates physical and psychosocial problems following ICU recovery</p> <ul style="list-style-type: none"> ● 4P: Patients, Physical, Psychosocial and Problems ● 4P comprises 53 items: 16 physical items, 26 psychosocial items and 11 follow-up ICU care items, scored on a 5-point Likert scale measuring level of agreement from “strongly agree” to “do not agree at all”
<p>5) Physical Function ICU Test (PFIT): used with critically ill patients who may not be able to mobilize away from the bedside, employing 4 domains</p> <ul style="list-style-type: none"> ● amount of assistance for sit to stand, rated from 0 (no physical assistance required) to 3 (assistance of 3 people required) ● strength for shoulder flexion and knee extension (rated on the Oxford Muscle Test Scale) ● marching in place (number of steps taken & the time required to complete these steps), & ● an upper extremity endurance task of arm elevation to 90° shoulder flexion (number of times both upper extremities are lifted above 90° of shoulder flexion)
<p>6) Karnofsky Performance Scale Index: a descriptive, ordinal scale that ranges from 100 (good health) to 0 (dead) and emphasizes physical performance and dependency</p> <ul style="list-style-type: none"> ● Karnofsky index of 70–100: a favorable functional outcome measure
<p>7) Modified Rankin Scale (mRS): quantifies independence and disability, with a scale of 6 grades (0-5)</p> <ul style="list-style-type: none"> ● 0, no symptoms; ● 1, no significant disability despite symptoms; ● 2, slight disability; ● 3, moderate disability; ● 4, moderately severe disability
<p>8) Glasgow Outcome Scale (GOS): provides a global assessment of function (see text for modified GOS scales)</p> <ul style="list-style-type: none"> ● score 1: good recovery; ● score 2: moderate disability; ● score 3: severe disability; ● score 4 vegetative state; ● score 5: death
<p>9) Disability Rating Scale (DRS): a common outcome measure of impairment, disability and handicap; the scale is intended to assess accurately general functional changes over the course of recovery</p> <ul style="list-style-type: none"> ● Impairment ratings: “Eye Opening”, “Communication Ability” and “Motor Response” ● Level of disability: Ability for “Feeding,” “Toileting” and “Grooming” ● Handicap: “Level of Functioning” and “Employability” ● Rating for each functioning area: scale of 0 to either 3 or 5 ● Maximum score (29): extreme vegetative state ● Lowest score (0): a person without disability

rehabilitation discharge and particularly at one year post injury, ceiling effects of the FIM were observed in the moderate and severely neurologically impaired population.²⁵

(C) FUNCTIONAL STATUS SCORE FOR THE ICU (FSS-ICU)

The FSS-ICU²¹ is an ordinal scale similar to FIM used for in-patient rehabilitation in ICU. The FSS-ICU consists of 3 pre-ambulation categories: (a) rolling, (b) supine to sit transfers and (3) unsupported sitting; and 2 ambulation categories: (a) sit to stand transfers and (b) ambulation. Each functional category is rated using a 1 (total dependent assistance) to 7 (complete independence) scale with a score from 0-35. A score of 0 is assigned if a patient is unable to perform a task due to physical limitations or medical status. The reliability and validity of the FSS-ICU has not been previously reported.¹⁰

(D) 4P QUESTIONNAIRE

Akerman et al²⁶ developed the 4P questionnaire for evaluating physical and psychosocial problems following ICU recovery. The questionnaire was named 4P after its major content: Patients, Physical, Psychosocial and Problems. It comprises 53 items: 16 physical items, 26 psychosocial items and 11 follow-up ICU care items. All items were based on a literature review and from clinical and theoretical experiences as ICU nurses regarding health and recovery after ICU²⁶.²⁴ The items were scored on a 5-point Likert scale measuring level of agreement from “strongly agree” to “do not agree at all”. There was also an option to answer “not relevant”. The questionnaire showed good construct validity in all three sets and had strong factor loadings for all three sets. The questionnaire has good concurrent validity compared with the Questionnaire SF-12. Internal consistency was shown to have reliable indices and stability reliability on retesting was good for the physical and psychosocial factors.²⁶

(E) PHYSICAL FUNCTION ICU TEST (PFIT)

The PFIT²⁷ is a reliable and responsive outcome measure that was developed for use with critically ill patients who may not be able to mobilize away from bedside. The test had 4 domains and has been shown to be both reliable and sensitive to change.²⁷ The 4 domains are: (a) amount of assistance for sit to stand, (b) strength for shoulder flexion and knee extension, (c) marching in place, and (d) an upper extremity endurance task of arm elevation to 90° shoulder flexion. The amount of assistance required to stand is rated from 0 (no physical assistance required) to 3 (assistance of 3 people required). Strength for shoulder flexion and knee extension is rated on the Oxford Muscle Test Scale. For marching in place, the examiner records the number of steps taken and the time required to complete these steps. For the upper extremity endurance component, the numbers of times both upper extremities are lifted above 90° of shoulder flexion are recorded as well as the time to complete this task. The PFIT has demonstrated reliability and

good responsiveness to change and thus may be advantageous to use as a supplement for documenting changes in mobility for a sub-population identified in the critical care environment.²⁷

(F) KARNOFSKY PERFORMANCE SCALE INDEX (KARNOFSKY STATUS SCALE/ KARNOFSKY INDEX/ KARNOFSKY SCORE)

The Karnofsky Index score²⁸ is among the recommended outcome measures for scoring of intensive care patients.¹³ It is used to give an indication of the patient’s functional status. This measurement tool is a well-established tool with proven validity and reliability for the assessment of independent functioning in the critically ill patients.¹³ It is a descriptive, ordinal scale that ranges from 100 (good health) to 0 (dead) and emphasizes physical performance and dependency. A Karnofsky index of 70–100 is generally considered a favorable functional outcome measure.

(G) MODIFIED RANKIN SCALE (MRS)

The mRS²⁹ quantifies independence and disability rather than performance of specific tasks. The scale consists of 6 grades from 0 to 5, as follows: 0, no symptoms; 1, no significant disability despite symptoms; 2, slight disability, whereby the subject is unable to carry out all previous activities, but is able to look after own affairs without assistance; 3, moderate disability when the subject requires some help, but is able to walk without assistance; 4, moderately severe disability with the subject being unable to attend to own bodily needs without assistance; 5, severe disability when the subject is bedridden, incontinent and requires constant nursing care and attention.³⁰ The scale was found to have good inter-rater agreement in acute stroke patients, but problems with the interpretation and relevancy of the scale in the hospitalized setting raise concerns about validity.³¹

(H) GLASGOW OUTCOME SCALE (GOS)

The GOS³² provides a global assessment of function and has been used in ICU settings.³³ It is well validated³⁴ and has a score of 1 which indicates a good recovery; 2 moderate disability; 3 severe disability; 4 vegetative state; and 5 death. In particular, the range of outcomes was described by classifying the patients as having minimal or no disability (Glasgow Outcome Score, 1), moderate disability (Glasgow Outcome Score 2), or severe disability (Glasgow Outcome Score, 3 or 4). Due to some shortcomings³⁵, the GOS was modified and a structured interview was proposed to accurately categorize patient’s disability. This version is called the GOS-Extended (GOSE/GOSe). This extended version of the scale separates each of the three higher function categories into two, thus making eight categories in total. It has been prospectively demonstrated that the validity criterion of the GOSe generally exceeds the GOS and it is more sensitive to change than the GOS. The GOSe has good reliability in neurological patients.^{36,37} Recently an alternate GOSE rating system has been proposed. First, patients are rated on GOS scores, and second, they are subcategorized on

GOSE scores, using structured questionnaire and narratives with central review committee. Through this system, GOSE scores can be categorized more accurately³⁸. The GOS includes good recovery and moderate disability and the GOSe includes severe disability, vegetative state and death.¹²

(I) DISABILITY RATING SCALE (DRS)

The DRS³⁹ is a common outcome measure of impairment, disability and handicap. The scale is intended to assess accurately general functional changes over the course of recovery. The first three items (“Eye Opening”, “Communication Ability” and “Motor Response”) reflect impairment ratings. Ability for “Feeding”, “Toileting” and “Grooming” reflect level of disability. The “Level of Functioning” and “Employability” reflects handicap. Each of the areas of functioning is rated on a scale of 0 to either 3 or 5, with the highest scores representing the higher level of disability. The maximum score is 29 (extreme vegetative state) and 0 for a person without disability. Many studies have reported good reliability and validity coefficients for the DRS.⁴⁰

DISCUSSION

Experimental studies have examined the functional ability of ICU patients during their stay and after discharge by using valid scales.⁷⁻¹⁰ However, no previous study has reviewed the use and psychometric properties of only functional assessment scales in a general ICU setting. Therefore, the purpose of the present review was to assess only the functional assessment scales for ICU patients, to examine reliability and validity of them, and to summarize their strengths and weaknesses.

Scales are usually the preferred method of assessing the functional outcome measure of patients. However, because scales are worded, scored, designed, and constructed differently, clinicians must be cautious when choosing an appropriate outcome measure. Different ICUs, such as stroke, cardiac, surgery, or respiratory units, determine specific functional outcome measures relevant to patient population. Outcome measures can also be classified according to their utility in specific settings, like acute stage, in-patient rehabilitation, and at follow up after discharge. Before choosing an outcome measure, one should first determine what needs to be measured, be it functional activity, impairment, capacity, performance, disability, and/or handicap and what type of administration one wants to be selected, e.g. testing in a laboratory, observation, or report by the patient. Each instrument should be reviewed for actual content and syllabus and instructions required for administration and scoring.¹²

Limitations still exist in all used scales when assessing patients in the ICU. First, there is a need for sufficient details in the questionnaire and in the instructions for the individual collecting the data to have reliable and valid responses. Second, a reasonable period of time is needed for improvement and stabilization of neurological, surgical or respiratory patients’

recovery before assessment. Therefore, a lot of useful time may be lost during the injury incidence and the functional measure which means drop-outs and loss of follow-up in clinical trials.¹¹

The outcome measures that have been reviewed in the present study appear to have specific shortcomings. Zanni et al²¹ reported that there are no methods for assessing functional status that have been specifically validated in ICU patients. Regarding the use of FIM for critical care environments, it is often employed in a subjective self-report format several months after ICU admission.⁴¹ Furthermore, several items of the total FIM cannot be assessed in an ICU setting (e.g., stairs) and a total score cannot be given. Although the ability of this tool to detect changes in the rehabilitation setting is high, there is a ceiling effect which limits the usefulness of it in assessing change after discharge from rehabilitation.²⁵ Also, it consolidates all transfer types (from bed mobility to standing transfers) into one task, which may lead to a floor effect for chronically critically ill populations.¹⁰ Gosselink et al⁴² report that the validated FIM does not evaluate basic mobility skills (e.g., rolling), which are more relevant for weak ICU patients. Until further validation work is done, this limitation is common to all publications in this field and was addressed through using methods similar to prior ICU publications.^{43,44}

However, the FSS-ICU includes tasks more appropriate for critically ill patients. The relevant to the ICU setting functional tasks, such as (1) rolling, (2) transfer from supine to sit, (3) sitting at the edge of bed, and (4) transfer from sit to stand, are evaluated. These 4 tasks, plus ambulation, were combined in the cumulative FSS-ICU, which is a simple sum of the 5 individual scores.²¹ However, reliability and validity of this new measure have not yet been reported.

Regarding the PFI test, although it is a reliable and responsive outcome measure, it is only described in a small sample of patients who were able to sit out of bed, stand from a chair, and march in place.²⁷ Many patients in ICU are not able to perform these out of bed tasks. Additionally, the PFIT does not assess ambulation. Therefore, the PFIT is likely to have both a floor and ceiling effect in an ICU population.¹⁰ Future research should aim to develop a PFIT score and investigate the ability of the PFIT to predict ICU readmission risk and functional outcome including ambulation.²⁷

The Karnofsky Index has been used to evaluate the 2–3 years of surviving re-admitted patients after discharge and identify possible determinants of their functional outcome.³³ Riachy et al⁴⁵ assessed the functional outcome of acute stroke patients at ICU discharge. The measurement properties of the Karnofsky Index have been subject to some limited investigation in critical care survivors. There is some evidence of their construct and criterion validity and their responsiveness, but reliability has not been investigated.¹³

The Barthel Index has been used widely, because it is short and it does not need experienced examiner. It has been used to measure physical functioning in clinical settings, especially neurological patients (i.e., stroke and traumatic brain injuries)

at ICU discharge^{7,46,47} and at baseline, discharge and follow up.⁴⁸ However, its psychological properties have received little attention, thus it is not possible to comment on their usefulness in critical care.²⁷ Therefore, due to limited research regarding its measurement properties in critical care patients and survivors, future studies should investigate the reliability, validity and responsiveness in the ICU environment.

The DRS is a common outcome measure of impairment, disability and handicap of neurological patients from the stage of coma in the ICU to their release to the community. Although the DRS is short and has an easy scoring system, it can assess general functioning rather than specific functional changes and cannot detect small functional changes in patients with mild impairment.⁴⁹ The major disadvantages of DRS relate to the fact that it requires more specialized training for its implementation by the rater, and to a high inter-rater variability.¹² The mRS quantifies independence and disability rather than performance of specific tasks particularly in stroke patients. Jeng et al⁴⁶ assessed the functional outcomes of 850 acute stroke patients at discharge from an ICU. Further studies should explore the Modified Ranking Scale's validity in the ICU setting.

The present study is an overview, as the authors have not conducted a systematic review and testing of the described assessment scales. Future studies may proceed to examine and test additional types of reliability and validity in all the existing functional scales with large ICU samples. Further investigations should focus on cross-cultural translation in the Greek language of the current instruments. The content validity must be assessed using a judge expert panel and the construct validity by using exploratory and confirmatory factor analysis. The exploratory factor analysis may examine the factor structure of the instrument. The confirmatory factor analysis further assesses the factorial validity supporting the instrument's model fit. Lastly, in future studies, it would also be beneficial to investigate whether a tool is used for research and/or for clinical purpose.

CONCLUSION

Assessing functional outcome is considered standard of care for the rehabilitation personnel, and is essential to document the effectiveness of treatment interventions. An important consideration when choosing a functional scale in the ICU is to determine first the patient population, its characteristics and its rehabilitation stage. Some measures may be most appropriate for individuals with severe activity limitations who are not able to sit, stand and walk, while other measures are most appropriate for patients who function at a higher level. The examiner may include a series of measures, some of which are not appropriate for patients at initial enrolment into a study, but are appropriate as the patient's condition improves minimal activity restrictions, and is ready for discharge to an assisted or independent living environment. To date only few outcome measures have been

developed specifically for chronically critically ill patients to measure function in a long-term acute care hospital setting. However, these current outcome measures have some limitations with their psychological properties. Therefore, further research should investigate the utility and the cross cultural validation/reliability of the existing functional outcome measures.

REFERENCES

1. Topp R, Ditmyer M, King K, Doherty K, Hornyak J III. The effect of bed rest and potential of prehabilitation on patients in the intensive care unit. *AACN Clin Issues* 2002; 13:263-276.
2. De Jonghe B, Lacherade JC, Durand MC, Sharshar T. Critical illness neuromuscular syndromes. *Crit Care Clin* 2007; 23:55-69.
3. Perme C, Chandrashekar R. Early mobility and walking program for patients in intensive care units: creating a standard of care. *Am J Crit Care* 2009; 18:212-221.
4. Morris PE. Moving our critically ill patients: mobility barriers and benefits. *Crit Care Clin* 2007; 23:1-20.
5. Needham DM. Mobilizing patients in the intensive care unit: improving neuromuscular weakness and physical function. *JAMA* 2008; 300:1685-1690.
6. Bruton A, Conway JH, Holgate ST. Reliability: what is it, and how is it measured? *Physiotherapy* 2000; 86:94-99.
7. van der Schaaf M, Dettling DS, Beelen A, et al. Poor functional status immediately after discharge from an intensive care unit. *Disabil Rehabil* 2008; 30:1812-1818.
8. Berney S, Haines K, Skinner E, Denehy L. Safety and feasibility of an exercise prescription approach to rehabilitation across the continuum of care for survivors of critical illness. *Phys Ther* 2012; 92:1524-1535.
9. Burtin C, Clerckx B, Robbeets C, et al. Early exercise in critically ill patients enhances short-term functional recovery. *Crit Care Med* 2009; 37:2499-2505.
10. Thrush A, Rozek M, Dekerlegand JL. The clinical utility of the Functional Status Score for the Intensive Care Unit (FSS-ICU) at a long-term acute care hospital: a prospective cohort study. *Phys Ther* 2012; 92:1536-1545.
11. Nichol AD, Higgins AM, Gabbe BJ, Murray LJ, Cooper DJ, Cameron PA. Measuring functional and quality of life outcomes following major head injury: Common scales and checklists. *Injury Int J Care Injured* 2011; 42:281-287.
12. Shuklaa D, Devia BI, Agrawalb A. Outcome measures for traumatic brain injury. *Clin Neurology Neuros* 2011; 113:435-441.
13. Black NA, Jenkinson C, Hayes JA, et al. Review of outcome measures used in adult critical care. *Crit Care Med* 2001; 29: 2119-2124.
14. Gosselink R, Needham D, Hermans G. ICU-based rehabilitation and its appropriate metrics. *Curr Opin Crit Care* 2012; 18: 533-539.
15. Mokkink LB, Terwee CB, Patrick DL, et al. International consensus on taxonomy, terminology, and definitions of measurement properties for health-related patient-reported outcomes: results of the COSMIN study. *J Clinical Epidem* 2010; 63:737-745.
16. Scholtes VA, Terwee CB, Poolman RW. What makes a meas-

- urement instrument valid and reliable? *Injury Int J Care Injured* 2011; 42:236–240.
17. Mahoney FI, Barthel DW. Functional evaluation: the Barthel Index. *MD State Med J* 1965; 14:61–65.
 18. Post MW, Visser-Meily JM, Gispens LS. Measuring nursing needs of stroke patients in clinical rehabilitation: A comparison of validity and sensitivity to change between the Northwick Park Dependency Score and the Barthel Index. *Clin Rehabil* 2002; 16:182–189.
 19. Granger CV, Hamilton BB, Keith RA, Zielesny M, Sherwin FS. Advances in functional assessment for medical rehabilitation. *Top Geriatr Rehabil* 1986; 1:59–74.
 20. Schweickert W, Pohlman M, Pohlman A, et al. Early physical and occupational therapy in mechanically ventilated, critically ill patients: a randomized controlled trial. *Lancet* 2009; 373:1874–1882.
 21. Zanni J, Korupolu R, Fan E, et al. Rehabilitation therapy and outcomes in acute respiratory failure: an observational pilot project. *J Crit Care* 2010; 25:254–262.
 22. Dodds TA, Martin DP, Stolov WC, Deyo RA. A validation of the Functional Independence Measurement and its performance among rehabilitation inpatients. *Arch Phys Med Rehabil* 1993; 74:531–536.
 23. Linacre JM, Heinemann AW, Wright BD, et al. The structure and stability of the Functional Independence Measure. *Arch Phys Med Rehabil* 1994; 75:127–132.
 24. Hamilton B, Laughlin J, Granger C, Kayton R. Interrater agreement of the seven level Functional Independence Measure (FIM). *Arch Phys Med Rehabil* 1991; 72:790.
 25. Hall KM, Mann N, High W, Wright J, Kreutzer J, Wood D. Functional measures after traumatic brain injury: ceiling effects of FIM, FIM + FAM, DRS and CIQ. *J Head Trauma Rehabil* 1996; 11:27–39.
 26. Akerman E, Fridlund B, Erssona A, Granberg-Axll A. Development of the 3-SET 4P questionnaire for evaluating former ICU patients' physical and psychosocial problems over time: A pilot study. *Inten Crit Care Nursing* 2009; 25:80–89.
 27. Skinner EH, Berney S, Warrillow S, Denehy L. Development of a physical function outcome measure (PFIT) and a pilot exercise training protocol for use in intensive care. *Critical Care and Resusc* 2009; 11:110–115.
 28. Schag CC, Heinrich RL, Ganz PA. Karnofsky performance status revisited: reliability, validity, and guidelines. *J Clin Oncol* 1984; 2:187–193.
 29. Van Swieten JC, Koudstaal PJ, Visser MC, Schouten HJA, van Gijn J. Inter observer agreement for the assessment of handicap in stroke patients. *Stroke* 1988; 19:604–607.
 30. Bonita R, Beaglehole R. Modification of Rankin Scale: recovery of motor function after stroke. *Stroke* 1988; 19:1497–1500.
 31. Zhao H, Collier JM, Quah DM, Purvis T, Bernhardt J. The Modified Rankin Scale in Acute Stroke Has Good Inter-Rater-Reliability but Questionable Validity. *Cerebrovasc Dis* 2010; 29:188–193.
 32. Jennett B, Bond M. Assessment of outcome after severe brain damage. *Lancet* 1975; 1:480–484.
 33. Conlon N, O'Brien B, Herbison GP, Marsh B. Long-term functional outcome and performance status after intensive care unit re-admission: a prospective survey. *British J Anaesth* 2008; 100:219–223.
 34. Kastebauer S, Pfister HW. Pneumococcal meningitis in adults: spectrum of complications and prognostic factors in a series of 87 cases. *Brain* 2003; 126:1015–1025.
 35. Anderson SI, Housley AM, Jones PA, Slattery J, Miller JD. Glasgow outcome score: an inter-rater reliability study. *Brain Inj* 1993; 7:309–317.
 36. Pettigrew LE, Wilson JT, Teasdale GM. Reliability of ratings on the Glasgow Outcome Scales from in-person and telephone structured interviews. *J Head Trauma Rehabil* 2003; 18:252–258.
 37. Wilson JT, Pettigrew LE, Teasdale GM. Structured interviews for the Glasgow Outcome Scale and the extended Glasgow Outcome Scale: guidelines for their use. *J Neurotrauma* 1998; 15:573–585.
 38. Lu J, Marmarou A, Lapane K, Turf E, Wilson L, et al. IMPACT Group. A method for reducing misclassification in the extended Glasgow outcome score. *J Neurotrauma* 2010; 27:843–852.
 39. Rappaport M, Hall KM, Hopkins K, et al. Disability Rating Scale for severe head trauma: coma to community. *Arch Phys Med Rehabil* 1982; 63:118–123.
 40. Eliason MR, Topp BW. Predictive validity of Rappaport's Disability rating scale in subjects with acute brain dysfunction. *Phys Ther* 1984; 64:1357–1360.
 41. Dennis DM, Hebdon-Todd TK, Marsh LJ, Cipriano LJ, Parsons RW. How do Australian ICU survivors fare functionally 6 months after admission? *Crit Care Resusc* 2011; 13:9–16.
 42. Gosselink R, Bott J, Johnson M, et al. Physiotherapy for adult patients with critical illness: recommendations of the European Respiratory Society and European Society of Intensive Care Medicine Task Force on Physiotherapy for Critically Ill Patients. *Inten Care Med* 2008; 34:1188–1199.
 43. De Jonghe B, Sharshar T, Lefaucheur JP, et al. Paresis acquired in the intensive care unit: a prospective multicenter study. *JAMA* 2002; 288:2859–2867.
 44. De Jonghe B, Bastuji-Garin S, Durand MC, et al. Respiratory weakness is associated with limb weakness and delayed weaning in critical illness. *Crit Care Med* 2007; 35:2007–2015.
 45. Riachy M, Sfeir F, Sleilaty G, et al. Prediction of the survival and functional ability of severe stroke patients after ICU therapeutic intervention. *BMC Neurology* 2008; 8:24.
 46. Jeng JS, Huang SJ, Tang SC, Yip PK. Predictors of survival and functional outcome in acute stroke patients admitted to the stroke intensive care unit. *J Neurol Sciences* 2008; 270:60–66.
 47. Navarrete-Navarro P, Rivera-Fernandez R, Lopez-Mutuberrria MT, et al. Outcome prediction in terms of functional disability and mortality at 1 year among ICU-admitted severe stroke patients: a prospective epidemiological study in the south of the European Union (Evascan Project, Andalusia, Spain). *Intensive Care Med* 2003; 29:1237–1244.
 48. Intiso D, Amoroso L, Zarrelli M, et al. Long-term functional outcome and health status of patients with critical illness poly-neuromyopathy. *Acta Neurol Scand* 2011; 123:211–219.
 49. Hall KM, Hamilton B, Gordon WA, Zasler ND. Characteristics and comparisons of functional assessment indices: disability rating scale. Functional independence measure and functional assessment measure. *J Head Trauma Rehabil* 1993; 8:60–74.